

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/363366437>

# Influence of Cactus Pear Mucilage-Based Edible Coating on Marketability and Edibility Parameters of Minimally Processed Loquat Fruits

Article in *Agronomy* · September 2022

DOI: 10.3390/agronomy12092120

CITATIONS

5

READS

38

6 authors, including:



**Giorgia Liguori**

Università degli Studi di Palermo

74 PUBLICATIONS 824 CITATIONS

[SEE PROFILE](#)



**Giuseppe Greco**

Università degli Studi di Palermo

6 PUBLICATIONS 26 CITATIONS

[SEE PROFILE](#)



**Raimondo Gaglio**

Università degli Studi di Palermo

131 PUBLICATIONS 2,024 CITATIONS

[SEE PROFILE](#)



**Luca Settanni**

Università degli Studi di Palermo

345 PUBLICATIONS 7,452 CITATIONS

[SEE PROFILE](#)

## Article

# Influence of Cactus Pear Mucilage-Based Edible Coating on Marketability and Edibility Parameters of Minimally Processed Loquat Fruits

Giorgia Liguori <sup>\*</sup>, Giuseppe Greco, Raimondo Gaglio , Luca Settanni , Paolo Inglese  and Alessio Allegra 

Department of Agricultural Food and Forest Sciences, University of Palermo, Viale delle Scienze, 90128 Palermo, Italy

\* Correspondence: giorgia.liguori@unipa.it

**Abstract:** Loquat fruit, popular all over the world for its mild, subacid, and sweet taste, has been classified as a non-climacteric fruit with a very short postharvest life. The fruit decays quickly after harvest, and losses in titratable acidity, taste, and juiciness, and internal browning occurs rapidly during shelf life. The aim of our study was to assess the effects of cactus pear mucilage-based coating on quality, nutraceutical value, microbiological growth, and sensorial parameters of minimally processed white-flesh *Martorana* and orange-red-flesh *Gigante Rossa* loquat fruits during cold storage. The effect of mucilage edible coating on the postharvest life, qualitative attributes, and nutraceutical value of fruit were evaluated by coloring, firmness, total soluble solids content, titratable acidity, ascorbic acid, antioxidant activity, total phenols, and total carotenoids content. Our data showed a significant effect of mucilage coating on preserving quality, nutraceutical value, sensorial parameters, and improving postharvest life of minimally processed loquat fruits. Furthermore, coated fruits showed a significantly lower microbiological growth than uncoated loquat fruits during the cold storage period. Our study suggests that minimally processing coated loquat fruit could allow producers to also sell to the market loquat fruits that present large spotted areas in the epicarp, which are usually considered unmarketable.

**Keywords:** *Eriobotrya japonica*; fresh-cut; antioxidant activity; microbiological growth



**Citation:** Liguori, G.; Greco, G.; Gaglio, R.; Settanni, L.; Inglese, P.; Allegra, A. Influence of Cactus Pear Mucilage-Based Edible Coating on Marketability and Edibility Parameters of Minimally Processed Loquat Fruits. *Agronomy* **2022**, *12*, 2120. <https://doi.org/10.3390/agronomy12092120>

Academic Editor: Pedro Javier Zapata

Received: 3 August 2022

Accepted: 1 September 2022

Published: 7 September 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Loquat (*Eriobotrya japonica* Lindl.), popular all over the world owing to the mild subacid, and sweet taste of its fruit, has been classified as a non-climacteric fruit with a very short postharvest life, with fruit decaying quickly after harvest, and losing titratable acidity, taste, and juiciness, and internal browning occurring rapidly during shelf life [1]. Chilling injury, browning, and purple spots are major problems, and the fruits are susceptible to various postharvest diseases, especially from mechanical injury and fruit processing [2].

Loquat cultivars show a wide diversity of fruit colors that is strictly related to the carotenoids content and distribution in fruit skin and flesh [3]. In addition to vitamins and minerals, loquat fruit is rich in phenolics and carotenoids. The bioactive components of loquat fruit include flavonoids, triterpenic acids, and carotenoids and show remarkably high scavenging activity against chemically generated radicals, thus making it effective in inhibiting oxidation of human low-density lipoproteins [2].

Loquat fruit suffer a quick senescence after harvest, and postharvest life is approximately 10 days in ambient conditions due to microbial decay, damage, and nutritional loss [4]. During postharvest storage, loquat fruit is affected by lignification, juiciness loss, internal browning, and microbial decay, which cause quality degradation and reduce its commodity value [4]. Several studies have reported that refrigeration has become one of the key postharvest factors and can combine with different treatments, as well as hot

air application, modified atmosphere, chemical and natural compounds, packaging, and edible coating to extend loquat fruits' postharvest life [1,3–5].

Edible coatings with semipermeable films can provide an alternative to a modified atmosphere—which is very expensive and not extensively used in commercial situations—by reducing cost, quality changes, and quantitative losses through the modification and control of the internal fruit atmosphere [3].

In recent years, significant changes in human lifestyles have produced an increase in popularity of fresh-cut foods that are ready to eat, and the consumption of minimally processed fruit and vegetables has undergone a sharp increase.

Cutting operations remove fruit epicarp that normally protects the internal tissue from microbiological invasion while allowing juices to leak from the inner tissues on the fruit surface. These effects of cutting explain the reason why microbial growth is much greater on a minimally processed product as compared with the corresponding intact product, and for that reason large microbiological populations, including potentially high levels of human pathogens, may develop on the surface of cut produce [6].

Among new postharvest management strategies of environment-friendly fresh fruit handling, the application of natural edible coatings has been reported to be very effective [7]. Edible coatings can act as a semipermeable barrier against gases and water vapor; can modify fruit tissue metabolism by affecting respiration rate, decreasing moisture and firmness loss, preserving the color, transporting antimicrobial, antioxidant, and other preservatives, controlling microbial growth, and maintaining fruit quality for a longer period [8].

A recent edible coating is the mucilage extracted from cladodes of *Opuntia ficus-indica*, which has been effectively used as a coating material, particularly for highly perishable fruits, minimally processed products, and fresh cut or sliced ones [7–14].

Studies have reported that *O. ficus-indica* edible coating positively affected fruit quality, reducing water transpiration and browning, maintaining fruit fresh weight, visual score values, fruit firmness, nutraceutical attributes, and controlling microbial growth, resulting in a longer storage period [8].

*O. ficus-indica* mucilage is a complex carbohydrate mixture composed of variable amounts of L-arabinose, D-galactose, L-rhamnose, and D-xylose, as well as galacturonic acid and quite large amounts of polyphenols, which could produce natural edible coating with a high nutraceutical value, useful for fruit and food preservation [7].

Despite several studies reporting the positive effect of edible coating when combined with low temperature and adequate humidity on handling and storage of loquat fruit, there is a lack of knowledge on the impact of coating treatments on the overall qualitative, sensorial, and nutraceutical value of minimally processed loquat fruits during cold storage. Few studies have reported results on postharvest performance of minimally processed loquat fruits [15], and no information exists on shelf life behavior of minimally processed loquat fruit treated with edible coatings.

Therefore, the aim of the present study was to evaluate the effect of the application of *O. ficus-indica* mucilage as an edible coating on pomological, physiochemical, sensorial and nutraceutical parameters, and microbial growth of minimally processed white-flesh *Martorana* and orange-red-flesh *Gigante Rossa* loquat cultivars during cold storage at  $5 \pm 0.5$  °C and 90% RH.

## 2. Materials and Methods

### 2.1. Loquat Fruit Samples

Loquat, *Eriobotrya japonica* Lindl, white-flesh (cv. *Martorana*—MRT) and orange-red-flesh (cv. *Gigante Rossa*—GR) fruits were harvested from commercial orchards located in Palermo (38°04' N, 13°23' E, 99–104 m a.s.l.). Loquat fruits were hand-picked at the ripe stage (light orange peel), suitable for the fresh fruit market and were quickly moved to the laboratory. Immediately after harvest, fruit quality parameters were analyzed using 30 fruits of each cultivar.

Loquat fruits were then washed under tap water, sanitized by immersion in 200 mg kg<sup>-1</sup> of sodium hypochlorite for 5 min, and left to dry at room temperature (~25 °C). Afterwards, fruits were peeled manually in a refrigerated room at 9 ± 1 °C. Only peeled fruits with no external injuries were selected, and fruit processing operations were carried out in sanitary conditions at 15 °C.

## 2.2. Fresh Mucilage Extraction and Application

One-year-old cladodes were collected from four-year-old *O. ficus-indica* (OFI) plants of the cultivar “Gialla”, located in the Department of Agricultural, Food and Forest Sciences, University of Palermo (38°7'4.0800" N 13°22'11.2800" E, 29 m a.s.l.). Cladodes (one-year-old cladodes) were harvested for mucilage extraction and moved to the laboratory where they were processed for mucilage extraction, using a modified patented method of Du Toit and De Witt developed in South Africa [8,16].

Cladodes were washed with chlorinated water to improve mucilage shelf life and to remove impurities and spines. Cladodes chlorenchyma was removed with a peeler to obtain pure high-quality mucilage. Cladodes were then sliced into squares and cooked in a microwave oven (900 W) for 3–5 min, until soft. The cooked soft cladode pieces were then mixed using an Omni Mixer Homogenizer (mod. Omni-Mixer. 17,107, Dupont Instruments Sorvall, Texas City, TX, USA) to aid the mucilage extraction. The obtained pulp was then centrifuged using a Sigma centrifuge (mod. 6K15, Sigma Laborzentrifugen GmbH, Osterode am Harz, Germany) at 8117× g for 15 min at 4 °C, to separate the liquid mucilage from the solids. The mucilage was then decanted and weighed while the solid material left in the falcon tubes was discarded. No chemicals were used during this extraction process and as such, the extracted mucilage obtained was natural and unadulterated by chemicals. Work surface area and cutting tools were washed and sanitized with 200 mg kg<sup>-1</sup> of sodium hypochlorite before and during fruit processing.

After cutting, loquat fruits were divided into two treatment groups for each cultivar (control: MRT and GR CTR and coated: MRT and GR OFI-EC). Each sampling group consisted in 5 replicates (3 fruits each) for each of 5 sampling dates, plus 25 replicates (3 fruits each) for sensory analysis and visual score (5 replicates for each sampling date) and 5 replicates (3 fruits each) for weight loss and respiration monitoring.

Loquat fruit OFI samples for each cultivar (MRT OFI-EC and GR OFI-EC) were treated with OFI mucilage, and loquat CTR samples (MRT CTR and GR CTR) were treated with distilled water and used as a control. Mucilage edible coating and distilled water were applied using an atomizing spray system (flow rate: 1 L h<sup>-1</sup>; air pressure: 50 kPa) [8,17]. Soon after coating, all fruits were air-dried at room temperature for 15' (~25 °C); then, coated and uncoated samples (OFI-EC and CTR), were placed in rigid polypropylene retail boxes 25 × 20 cm retail boxes (3 peeled loquat fruits to each box), sealed with 35 µm microperforated polypropylene film (O<sub>2</sub> permeability: ~12,000 mL m<sup>-2</sup> d<sup>-1</sup> atm<sup>-1</sup>; CO<sub>2</sub> permeability: ~13,000 mL m<sup>-2</sup> d<sup>-1</sup> atm<sup>-1</sup> at 5 °C) and stored at 5 ± 0.5 °C and 95% RH for 13 days.

## 2.3. Quality Parameters: Firmness, Soluble Solid Content, Titratable Acidity, Color, and Weight Loss

The quality of white and orange-red-flesh minimally processed loquat fruits was assessed soon after coating (0 d) and at 3, 5, 7, 10, and 13 days of storage at 5 °C. For each sampling date and experimental treatment, five samples (3 fruits for each) of loquat fruits were randomly chosen and analyzed.

Fruit firmness was measured using a digital penetrometer (TR model 53,205, Turoni, Forlì, Italy) incorporating an 8 mm-diameter probe, after removal of a small piece of peel.

Average values were calculated from the results of 15 fruits' measurements (2 measurements per fruit) for each treatment and cultivar at each sampling date.

Extractable juice was determined by the weight loss from tissue disks (after centrifuging for 10 min at  $1500 \times g$  at ambient temperature [4]). The results of extractable juice were expressed as the percentage fresh weight loss of the tissue plugs after centrifugation.

After firmness determinations, fruit pulp was homogenized and used to measure total soluble solids (TSS) content and titratable acidity (TA). Total soluble solids content (TSS) was determined by a digital refractometer (Palette PR-32, Atago Co., Ltd., Tokyo, Japan); titratable acidity (TA) was measured by titration of 10 mL homogenized fruit flesh juice with 0.1 N NaOH to an endpoint of pH 8.1 and expressed as the percentage of citric acid (mod. S compact titrator, Crison Instruments, Barcelona, Spain).

Weight loss for each loquat fruit cultivar was calculated using 5 packages for each treatment (5 boxes of 3 fruits  $\times$  2 treatments per cultivar) and expressed as the percentage reduction with respect to the initial time, using a two-decimal precision digital balance (mod. CENT-2 10000, Gibertini, Milan, Italy).

$$\% \text{ Weight loss} = [(W_i - W_s)]/W_i \times 100$$

where  $W_i$  is the initial weight, and  $W_s$  is the weight measured during storage.

Minimally processed loquat fruits' external color for each cultivar and treatment were measured at two opposite points on each fruit using a colorimeter (Chroma Meter CR-400 C, Minolta, Osaka, Japan). CIE  $L^*a^*b^*$  coordinates were recorded as  $L^*$  (lightness),  $a^*$  (positive values for reddish colors and negative values for greenish colors), and  $b^*$  (positive values for yellowish colors and negative values for bluish colors).

#### 2.4. Headspace Gas Composition

In packages,  $O_2$  and  $CO_2$  partial pressure were measured immediately before quality evaluation, using an  $O_2$  and  $CO_2$  portable analyzer (Dansensor Checkpoint, Mocon, Minneapolis, MN, USA) after 0, 3, 5, 7, 10, and 13 days at  $5^\circ C$  using 5 packages for each cultivar and treatment.

#### 2.5. Nutraceutical Attributes

Total phenolic content, carotenoids, ascorbic acid content, and antioxidant activity of minimally processed loquat fruits of each cultivar were assessed soon after coating (0 d) and at 3, 5, 7, 10, and 13 days of storage at  $5^\circ C$ . For each sampling date and experimental treatment (MRT-CTR, GR-CTR, and MRT OFI-EC, GR OFI-EC), three samples were randomly chosen and analyzed.

##### 2.5.1. Fruit Extract Preparation

Loquat fruit samples were frozen at  $-80^\circ C$  until extract preparation. The frozen samples were thawed and chopped and the seeds were separated from the pulp. The pulp was homogenized, and fruit extracts were prepared as previously described with minor changes [18]. Briefly, ten grams of the whole homogenate was weighed and then extracted with MeOH using a 1:5 ( $w/v$ ) ratio. Samples were mixed by vortex for 5 min and sonicated at room temperature for 15 min ( $\sim 25^\circ C$ ). The mixtures were allowed to stand for 2 h at room temperature. After centrifugation (10 min at  $8000 \times g$ ,  $4^\circ C$ ) the supernatants were filtered, portioned, and stored at  $-20^\circ C$ . The extraction procedure was repeated to obtain three different technical replicates.

##### 2.5.2. Total Phenolic and Total Carotenoids Content

Total phenolic content (TPC) of extracts from treated and untreated fruits of each loquat cultivar was determined by the reduction of phosphotungstic-phosphomolybdic acid (Folin-Ciocalteu's reagent) to blue pigments, in alkaline solution according to Folin-Ciocalteu's method [19]. Total phenolic content was expressed as mg gallic acid (GA) equivalent (GAE).

The total carotenoids were extracted in flesh loquat fruits as described by Petriccione et al. [3] and determined spectrophotometrically according to Kichtenthaler and

Wellburn [20]. Results were expressed as milligrams per 100 g fresh weight (FW), all measurements were done in three replicates.

### 2.5.3. DPPH and Hydroxyl Radical Scavenging Analysis

The 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging analysis was performed on 2 g of frozen tissue with 5 mL 50% (*v/v*) ethanol according to Wang et al. [4]. The homogenate was centrifuged at  $12,000\times g$  for 20 min at 4 °C, then the supernatant was collected. The reaction mixture consisted of 0.1 mL supernatant and 1.9 mL  $120\ \mu\text{mol L}^{-1}$  DPPH. Results were calculated using the following formula: DPPH radical scavenging activity (%) =  $[(A_0 - A_1)/A_0] \times 100$ , with  $A_0$  referring to absorbance of the control,  $A_1$  referring to absorbance of the samples. DPPH radical scavenging was expressed as a percentage.

The hydroxyl radical scavenging analysis was performed on 2 g of frozen tissue with 5 mL 50% (*v/v*) ethanol according to Wang et al. [4]. The homogenate was centrifuged at  $12,000\times g$  for 20 min at 4 °C, then the supernatant was collected. The reaction mixture consisted of 0.5 mL supernatant, 1.5 mL salicylic acid, 2 mL water, and 0.1 mL 0.3%  $\text{H}_2\text{O}_2$ . Results were calculated following the formula: hydroxyl radical scavenging (%) =  $[(A_0 - A_1)/A_0] \times 100$ , with  $A_0$  referring to absorbance of the control,  $A_1$  referring to absorbance of the samples. The hydroxyl radical scavenging was expressed as a percentage.

### 2.5.4. Ascorbic Acid Content

Ascorbic acid in MRT and GR- CTR and MRT and GR-OFI samples was determined by extracting 10 g of blended fruit sample in 100 mL metaphosphoric acid ( $\text{HPO}_3$ ), then filtering it through Whatman no. 1 filter paper. A volume of 10 mL filtered solution was determined volumetrically with the 2–6 dichlorophenol-indophenol reagent until a slightly pink coloration was observed and persisted for 15 s [21]. The ascorbic acid content was expressed in mg/100 g FW.

## 2.6. Sensory Analysis and Visual Score

On each sampling date, 5 boxes (3 fruits in each) for each treatment (MRT and GR-CTR, and MRT and GR-OFI) were subjected to sensory evaluation. The sensory profile was constructed by a panel of 10 judges trained in a few preliminary meetings: by using commercial fruit, the judges generated a list of descriptors. Sensory analysis was focused on appearance, sweetness, acidity, aroma, off-flavor development, taste, texture, juiciness, and overall acceptance. The different descriptors were quantified using a ten-point intensity scale where the digit 1 indicates the descriptor absence while the digit 10 the full intensity [7,8]. The order of presentation was randomized between judges. Water was provided for rinsing between samples.

On each sampling date, 5 boxes (3 fruits in each) for each treatment (MRT and GR-CTR, and MRT and GR- OFI) were also evaluated by each judge for the visual score. Visual appearance score resulted from the medium value of color, visible structural integrity, and visual appearance [8]. The different descriptors were quantified using a subjective 5–1 rating scale with 5 = very good, 4 = good, 3 = sufficient, 2 = poor (limit of edibility), and 1 = very poor (inedible) [8]. A score of 3 was the limit of marketability. The order of presentation was randomized between judges.

## 2.7. Microbiological Characterization of *O. ficus-indica* Mucilage Edible Coating

### 2.7.1. Determination of Antibacterial Activity

*O. ficus-indica* mucilage edible coating was tested against several bacterial strains of food origin representing spoilage and pathogenic bacterial groups. In particular, *Pseudomonas endophytica* and *P. fluorescens* were chosen among spoilage bacteria, while *Escherichia coli*, *Listeria monocytogenes*, *Salmonella enteritidis*, and *Staphylococcus aureus* were among the main food-borne bacterial pathogens. All indicator strains were reactivated in Brain Heart Infusion (BHI) broth (Condalab, Madrid, Spain). The incubation was carried out for 24 h

at 37 °C, excepting *Pseudomonas* strains, which were incubated at 25 °C. The antibacterial activity of *O. ficus-indica* mucilage edible coating was tested by the paper disc diffusion (PDD) method [22] using sterile water as a negative control, while streptomycin (10% *w/v*) was a positive control [23]. The inhibitory activity was evaluated after 24 h of incubation at 25 °C and 37 °C for spoilage and pathogenic bacteria, respectively, and was scored positive only in case of a definite clear area around the paper discs. The tests were carried out in triplicate.

#### 2.7.2. Plate Count

One milliliter of *O. ficus-indica* mucilage edible coating was directly subjected to decimal serial dilutions (1:10) in Ringer's solution (Sigma-Aldrich, Milan, Italy) and the microbial suspensions were used for the search of the main undesired communities, including both spoilage and pathogenic populations. The inoculation, cultivation, and incubation of the different microbial groups occurred as described by Passafiume et al. [17]. Briefly, total mesophilic microorganisms (TMM) and total psychotrophic microorganisms (TPM) were detected on plate count agar (PCA), *Pseudomonas* spp. on *Pseudomonas* agar base (PAB) with added fucidin cephaloridine supplement (CFC), members of the Enterobacteriaceae family on violet red bile glucose agar (VRBGA), coagulase-positive staphylococci (CPS) on Baird Parker (BP) supplemented with rabbit plasma fibrinogen (RPF), *Listeria monocytogenes* on *Listeria* selective agar base with SR0140E supplement, *Salmonella* spp. and *Escherichia coli* on Hektoen enteric agar (HEA), and yeasts and molds on yeast extract peptone dextrose (YPD) agar supplemented with chloramphenicol (0.1 g/L) to prevent bacterial growth. All media and chemicals were purchased from Microbiol Diagnostici (Uta, Italy). Plate counts were performed in triplicate.

#### 2.8. Microbiological Analyses of Loquat Fruit

Coated and untreated loquat samples (25 g) from each cultivar were homogenized in 225 mL of Ringer's solution (Sigma-Aldrich) by a stomacher Bag-Mixer 400 (Interscience, Saint Nom, France) for 3 min at the highest speed (blending power 4). Homogenized samples were then serially diluted. *O. ficus-indica* mucilage edible coating was diluted as reported above and the cell suspensions were subjected to the enumeration of spoilage and pathogenic microorganisms. Plate counts were performed in triplicate at each collection time (soon after production and after 3, 5, 7, 10, and 13 days of refrigerated storage).

#### 2.9. Statistical Analyses

All data were submitted to one-way analysis of variance (ANOVA) and means were separated with Tukey's test at  $p \leq 0.05$ . The statistical analysis was carried out using Systat 10 (Systat, Chicago, IL, USA).

### 3. Results

#### 3.1. Quality Parameters: Firmness, Soluble Solid Content, Titratable Acidity, Extractable Juice, Ascorbic Acid Content, Color, and Weight Loss

Fruit firmness decreased significantly in *Martorana* (MRT) and *Gigante Rossa* (GR) cultivars during storage in both treatments (Table 1). Significant differences between MRT CTR and MRT OFI-EC and GR CTR and GR OFI-EC samples occurred from the third day of storage at 5 °C until the end of the storage (Table 1). MRT CTR and GR CTR samples showed the highest decreases in firmness of 41% and 29%, respectively, from  $T_0$  to the end of the cold storage period (Table 1). Otherwise, MR OFI-EC and GR OFI-EC showed the highest fruit firmness values at the end of the storage with losses of firmness of 33% and 24%, respectively, from the beginning to the end of the cold storage period, showing the effectiveness of OFI-EC mucilage coating in terms of maintaining fruit cell structure (Table 1). GR samples showed the lowest loss of firmness in both treatments compared to MRT ones (Table 1). Concerning chemical parameters, there was evidence of slight changes in terms of TSS and TA during storage in both MRT and GR CTR and MRT and GR OFI-EC

samples (Table 1). TSS showed a slight decrease in MRT CTR and MRT OFI-EC samples during cold storage, with no significant differences between the treatments until the tenth day of cold storage; significant differences between the treatments were shown only at the end of the cold storage period (13 days at 5 °C) (Table 1). GR OFI-EC showed significantly higher TSS values on all the sampling dates compared to GR CTR ones (Table 1).

**Table 1.** Changes in firmness, total soluble solids (TSS) titratable acidity (TA), extractable juice, and ascorbic acid content in minimally processed loquat fruits (cv *Martorana* and cv *Gigante Rossa*), untreated (CTR) samples, and those treated with mucilage (OFI-EC) during cold storage (13 days at 5 °C). Different lowercase letters indicate significant differences at  $p \leq 0.05$  between the treatments on each sampling date. Data are the mean  $\pm$  SE ( $n = 5$ ).

Variety	Storage Time (days)	Treatments	Firmness (N)	TSS (°Brix)	TA (% malic acid)	Extractable Juice (%)	Ascorbic Acid (mg 100 g <sup>-1</sup> FW)
<i>Martorana</i>	0	CTR	8.49 $\pm$ 0.12	-	10.15 $\pm$ 0.34	-	6.14 $\pm$ 0.03
	3	CTR	7.75 $\pm$ 0.09	<i>b</i>	10.01 $\pm$ 0.23	<i>b</i>	6.14 $\pm$ 0.03
	5	CTR	7.35 $\pm$ 0.08	<i>b</i>	9.51 $\pm$ 0.41	<i>b</i>	4.95 $\pm$ 0.05
	7	CTR	6.72 $\pm$ 0.14	<i>b</i>	9.37 $\pm$ 0.21	<i>b</i>	4.95 $\pm$ 0.05
	10	CTR	6.10 $\pm$ 0.02	<i>b</i>	9.18 $\pm$ 0.29	<i>b</i>	4.95 $\pm$ 0.05
<i>Martorana</i>	0	OFI-EC	8.49 $\pm$ 0.12	-	10.15 $\pm$ 0.34	-	6.14 $\pm$ 0.03
	3	OFI-EC	8.25 $\pm$ 0.08	<i>a</i>	10.05 $\pm$ 0.29	<i>a</i>	6.14 $\pm$ 0.03
	5	OFI-EC	7.81 $\pm$ 0.09	<i>a</i>	9.58 $\pm$ 0.32	<i>a</i>	5.15 $\pm$ 0.01
	7	OFI-EC	7.19 $\pm$ 0.11	<i>a</i>	9.48 $\pm$ 0.29	<i>a</i>	5.15 $\pm$ 0.01
	10	OFI-EC	6.91 $\pm$ 0.03	<i>a</i>	9.17 $\pm$ 0.14	<i>a</i>	5.15 $\pm$ 0.01
<i>Gigante Rossa</i>	0	CTR	10.94 $\pm$ 0.32	-	10.33 $\pm$ 0.29	-	5.13 $\pm$ 0.04
	3	CTR	8.92 $\pm$ 0.19	<i>b</i>	9.23 $\pm$ 0.13	<i>b</i>	3.12 $\pm$ 0.02
	5	CTR	8.74 $\pm$ 0.12	<i>b</i>	9.08 $\pm$ 0.21	<i>b</i>	3.12 $\pm$ 0.02
	7	CTR	8.53 $\pm$ 0.08	<i>b</i>	8.68 $\pm$ 0.24	<i>b</i>	3.12 $\pm$ 0.02
	10	CTR	8.31 $\pm$ 0.10	<i>b</i>	8.47 $\pm$ 0.14	<i>b</i>	3.12 $\pm$ 0.02
<i>Gigante Rossa</i>	0	OFI-EC	10.94 $\pm$ 0.32	-	10.33 $\pm$ 0.29	-	5.13 $\pm$ 0.04
	3	OFI-EC	9.88 $\pm$ 0.12	<i>a</i>	9.97 $\pm$ 0.61	<i>a</i>	5.13 $\pm$ 0.04
	5	OFI-EC	9.39 $\pm$ 0.09	<i>a</i>	9.75 $\pm$ 0.18	<i>a</i>	4.97 $\pm$ 0.04
	7	OFI-EC	9.15 $\pm$ 0.11	<i>a</i>	9.60 $\pm$ 0.19	<i>a</i>	4.97 $\pm$ 0.04
	10	OFI-EC	8.92 $\pm$ 0.09	<i>a</i>	9.23 $\pm$ 0.11	<i>a</i>	4.97 $\pm$ 0.04
<i>Gigante Rossa</i>	0	OFI-EC	8.33 $\pm$ 0.07	<i>a</i>	9.21 $\pm$ 0.08	<i>a</i>	3.84 $\pm$ 0.03
	3	OFI-EC	8.33 $\pm$ 0.07	<i>a</i>	9.21 $\pm$ 0.08	<i>a</i>	3.84 $\pm$ 0.03
	5	OFI-EC	8.33 $\pm$ 0.07	<i>a</i>	9.21 $\pm$ 0.08	<i>a</i>	3.84 $\pm$ 0.03
	7	OFI-EC	8.33 $\pm$ 0.07	<i>a</i>	9.21 $\pm$ 0.08	<i>a</i>	3.84 $\pm$ 0.03
	10	OFI-EC	8.33 $\pm$ 0.07	<i>a</i>	9.21 $\pm$ 0.08	<i>a</i>	3.84 $\pm$ 0.03

TA values showed a slight decrease during cold storage in both MRT and GR CTR and MRT and GR OFI-EC samples, showing higher values in MRT OFI-EC and GR OFI-EC samples than those in MRT CTR and GR CTR ones until the end of the cold storage period (Table 1).


OFI-EC edible coating maintained a higher level of extractable juice than control fruits of both cultivars (Table 1). The extractable juice of MRT OFI-EC and GR OFI-EC samples was 4.78% and 3.78% greater than that of MRT CTR and GR CTR ones at the end of the cold storage period (13 days at 5 °C), respectively (Table 1).

Higher content of ascorbic acid was found in MRT OFI-EC and GR OFI-EC samples than MRT CTR and GR CTR ones until the end of the cold storage period (Table 1). Significantly, ascorbic acid content of OFI-EC coated samples of cultivars was 10.54% and 9.37% higher, respectively, than that of control samples at the end of the cold storage period (Table 1). Therefore, OFI edible coating significantly inhibited the decrease of firmness and TSS and maintained higher levels of extractable juice and ascorbic acid after 13 days of storage at 5 °C in both loquat fruit cultivars.


The loss of weight progressively increased with storage time and was linear for both cultivars and treatments. The mucilage coating treatment significantly decreased weight loss percentage during cold storage in MRT OFI-EC and GR OFI-EC samples compared



to that of MRT CTR and GR CTR ones (Figures 1 and 2). Indeed, MRT CTR and GR CTR samples showed a weight loss 1.5 times higher than that of MRT OFI-EC and GR OFI-EC samples at the end of the cold storage (Figures 1 and 2). Differences between coated and uncoated fruit were significant starting from 2 days until the end of the cold storage period in both cultivars (Figures 1 and 2). OFI edible coating was significantly effective in preventing weight loss in both cultivars.




**Figure 1.** Changes in weight loss (%) in minimally processed *Martorana* loquat fruits, untreated fruit (MRT CTR), and fruits treated with mucilage (MRT OFI-EC) during cold storage (13 days at 5 °C). Different lowercase letters indicate significant differences at  $p \leq 0.05$  between the treatments on each sampling date. Data are the mean  $\pm$  SE (Vertical bars represent standard error;  $n = 5$ ).



**Figure 2.** Changes in weight loss (%) in minimally processed *Gigante Rossa* loquat fruits, untreated fruit (GR CTR), and fruits treated with mucilage (GR OFI-EC) during cold storage (13 days at 5 °C). Different lowercase letters indicate significant differences at  $p \leq 0.05$  between the treatments on each sampling date. Data are the mean  $\pm$  SE (Vertical bars represent standard error;  $n = 5$ ).

Fruit flesh brightness ( $L^*$ ) was similar in loquat fruits of both cultivars at the time of treatment. Fruit flesh color slightly decreased in *Martorana* (MRT) and *Gigante Rossa* (GR) cultivars during storage in both treatments (Figure 3).




**Figure 3.** Changes in brightness ( $L^*$ ) in minimally processed loquat (cv *Martorana* and *Gigante Rossa*), untreated fruit (CTR), and fruits treated with mucilage (OFI-EC) during cold storage (13 days at 5 °C). Data are the mean  $\pm$  SE (bars represent standard error of the means;  $n = 5$ ).

MRT CTR and MRT OFI-EC samples showed similar values until 3 days of cold storage and then MRT CTR samples showed lower values of flesh brightness than those of MRT OFI-EC ones until the end of the cold storage period (from 3 to 13 days of storage at 5 °C) (Figure 3). MRT CTR and MRT OFI-EC showed a slight decrease during storage, with a loss of 10% and 13% of flesh brightness, respectively, from T0 to 13 days of cold storage (Figure 3).

GR CTR and GR OFI-EC samples showed a sharp decrease from T0 to 3 days of cold storage, then flesh brightness maintained similar values until 10 days of cold storage in both treatments (Figure 2), while at the end of the cold storage period, GR CTR flesh brightness decrease was 2 times higher than that in GR OFI-EC samples (Figure 3). The mucilage coating positively affected fruit quality parameters, reduced weight loss, and improved fruit brightness.

### 3.2. Headspace Gas Composition

In-package atmosphere was significantly affected by storage time in both treatments and cultivars. During cold storage, a decrease in  $O_2$  and an increase in  $CO_2$  in-package levels were observed for MRT CTR, MRT OFI-EC, GR CTR, and GR OFI-EC packaging (Figure 4A,B). MRT CTR and GR CTR samples showed a significantly higher level of  $CO_2$  than that of MRT OFI-EC and GR OFI-EC during storage; indeed, MRT CTR and GR CTR samples showed an in-package  $CO_2$  concentration almost twice that in MRT OFI-EC and GR OFI-EC ones after 13 days of cold storage (Figure 4A). MRT OFI-EC and GR OFI-EC samples showed a significantly higher level of  $O_2$  than that of MRT CTR and GR CTR during storage, with values 1.5 times higher than that in MRT CTR and GR CTR samples at the end of the cold storage period (Figure 4B).



**Figure 4.** Concentrations of CO<sub>2</sub> (A) and O<sub>2</sub> (B) in minimally processed loquat (cv *Martorana* and *Gigante Rossa*), untreated fruit (CTR), and fruits treated with mucilage (OFI-EC) during cold storage (13 days at 5 °C). Data are the mean  $\pm$  SE (bars represent standard error of the means;  $n = 5$ ).

After 13 days of cold storage, the O<sub>2</sub>/CO<sub>2</sub> in-package concentration (kPa) in MRT CTR and MRT OFI-EC, and GR OFI-EC and GR CTR was about 6/12, 9/5, 7/8, and 11/4, respectively (Figure 4A,B). MRT CTR and GR CTR fruits showed a higher respiration rate during cold storage than that of MRT OFI-EC and GR OFI-EC fruits; indeed, MRT CTR samples showed a loss in terms of in-package O<sub>2</sub> concentration of 74% from the beginning to the end of the cold storage; otherwise, the in-package O<sub>2</sub> concentration loss in MRT OFI-EC was 54% from the beginning to the end of the cold storage period (Figure 4B). Similar behavior in GR CTR samples showed a loss in terms of in-package O<sub>2</sub> concentration of 64% from the beginning to the end of the cold storage period, while the in-package O<sub>2</sub> concentration loss in GR OFI-EC was 44% from the beginning to the end of the cold storage period (Figure 4B).

### 3.3. Bioactive Compounds (Total Phenolic and Total Carotenoids Content) and Radical Scavenging Activity

Total phenolic content was significantly affected by storage time and treatment (Table 2). Total phenolic content in MRT and GR showed an increase at day 5, then de-

creased until the end of the cold storage period in both treatments (Table 2). Total phenolic content in *Martorana* loquat fruits was noticeably higher than that in *Gigante Rossa* ones (Table 2). OFI edible coating in both cultivars induced the increase and prevented the decline of total phenolic content. Phenolic content was 14% and 27% higher, respectively, in MRT OFI-EC and GR OFI-EC, than that in MRT CTR and GR CTR ones at the end of the cold storage period (Table 2).

**Table 2.** Total phenolic and total carotenoids content in minimally processed loquat fruits (cv *Martorana* and cv *Gigante Rossa*), untreated fruit (CTR), and fruits treated with mucilage (OFI-EC) during cold storage (13 days at 5 °C). Different lowercase letters indicate significant differences at  $p \leq 0.05$  between the treatments on each sampling date. Data are the mean  $\pm$  SE ( $n = 5$ ).


Variety	Storage Time (days)	Treatments	Total Phenolic Content (mg GAE/100 g FW)		Total Carotenoids (mg/100 g FW)	
<i>Martorana</i>	0	CTR	48.24 $\pm$ 0.89	-	1.30 $\pm$ 0.03	-
	5	CTR	54.89 $\pm$ 1.04	<i>b</i>	1.40 $\pm$ 0.07	-
	13	CTR	50.41 $\pm$ 0.95	<i>b</i>	1.65 $\pm$ 0.08	-
<i>Martorana</i>	0	OFI-EC	48.24 $\pm$ 0.89	-	1.30 $\pm$ 0.03	-
	5	OFI-EC	61.35 $\pm$ 1.09	<i>a</i>	1.45 $\pm$ 0.05	-
	13	OFI-EC	57.23 $\pm$ 0.56	<i>a</i>	1.60 $\pm$ 0.04	-
<i>Gigante Rossa</i>	0	CTR	29.03 $\pm$ 0.43	-	1.50 $\pm$ 0.06	-
	5	CTR	38.41 $\pm$ 0.38	<i>b</i>	1.90 $\pm$ 0.04	-
	13	CTR	30.15 $\pm$ 0.52	<i>b</i>	2.08 $\pm$ 0.03	-
<i>Gigante Rossa</i>	0	OFI-EC	29.03 $\pm$ 0.43	-	1.50 $\pm$ 0.06	-
	5	OFI-EC	49.36 $\pm$ 0.41	<i>a</i>	1.85 $\pm$ 0.04	-
	13	OFI-EC	38.42 $\pm$ 0.38	<i>a</i>	2.00 $\pm$ 0.07	-

Total carotenoids content showed a slight increase during the cold storage in both treatments and cultivars (Table 2). Total carotenoids content in *Gigante Rossa* loquat fruits was slightly higher than that in *Martorana* ones (Table 2). OFI edible coating treatment did not affect the total carotenoids content in either cultivar.

The radical scavenging activity (DPPH) increased until the fifth day of cold storage and then decreased until the end of the cold storage period in both treatments and cultivars (Figure 5A). The DPPH radical scavenging rate in *Martorana* fruits was higher than that in *Gigante Rossa* loquat fruits (Figure 5A). DPPH radical scavenging rate in MRT OFI-EC and GR OFI-EC samples was significantly higher than that in MRT CTR and GR CTR ones, showing values 32% and 36% higher, respectively, at the end of the cold storage period (Figure 5A).

Hydroxyl radical scavenging rate increased until the fifth day of cold storage and then maintained similar values until the end of the cold storage period in both treatments and cultivars (Figure 5B). MRT OFI-EC and GR OFI-EC samples showed values significantly higher than those in MRT CTR and GR CTR ones; the hydroxyl radical scavenging rate was 20%, 22%, 4%, and 3% higher from T0 until the end of the cold storage period, respectively (Figure 5B).

OFI edible coating significantly enhanced the hydroxyl radical scavenging rate in loquat fruits and maintained a higher hydroxyl radical scavenging rate in comparison with the control MRT and GR cultivars until the end of the cold storage period.



**Figure 5.** DPPH radical scavenging rate (A) and hydroxyl radical scavenging rate (B) in minimally processed loquat (cv *Martorana* and *Gigante Rossa*), untreated fruit (CTR), and fruits treated with mucilage (OFI-EC) during cold storage (13 days at 5 °C). Data are the mean  $\pm$  SE (bars represent standard error of the means;  $n = 5$ ).

### 3.4. Inhibitory Properties and Microbiological Characterization of *O. ficus-indica* Mucilage Edible Coating

The antibacterial activity of *O. ficus-indica* mucilage against spoilage and pathogenic bacteria is shown in Table 3. Except for the strain *E. coli* ATCC25922, all other indicator strains were inhibited by the edible coating with a diameter of the inhibition area around the paper discs higher than 12 mm.

The microbiological counts of *O. ficus-indica* mucilage edible coating did not reveal the presence of spoilage and pathogenic microbial populations, showing its high hygienic suitability for food production.

**Table 3.** Antibacterial activity of *O. ficus-indica* mucilage edible coating.

Species	Strains	Source of Isolation	Inhibition (mm)
Spoilage			
<i>Pseudomonas endophytica</i>	4G764	Ready to eat salad	14.8 ± 0.2
<i>Pseudomonas fluorescens</i>	4G628	Ready to eat salad	13.9 ± 0.2
Pathogenic			
<i>Escherichia coli</i>	ATCC25922	Clinical isolate	-
<i>Listeria monocytogenes</i>	ATCC19114	Animal tissue	16.3 ± 0.3
<i>Salmonella enteritidis</i>	ATCC13076	Unknown	13.7 ± 0.1
<i>Staphylococcus aureus</i>	ATCC33862	Unknown	12.1 ± 0.2

Results indicate the mean value of three independent assays. Symbols: - no inhibition found.

### 3.5. Evolution of Microbiological Parameters on Loquat Fruits

The results of plate counts performed on untreated (MRT-CTR and GR-CT) and coated (MRT OFI-EC and GR OFI-EC) loquat samples of both cultivars that were objects of study (*Martorana* and *Gigante Rossa*) during refrigerated storage are reported in Table 4. The presence of members of the Enterobacteriaceae family, CPS, *Salmonella* spp., and *L. monocytogenes* was not found (for this reason, these results are not reported in Table 4).

**Table 4.** Microbial loads of untreated and coated loquat samples.

Storage Time	Samples	Microbial Loads				
		TMM	TPM	Pseudomonads	Yeasts	Molds
0 days	MRT CTR	<2 a	<2 a	<2 a	<2 a	<2 a
	MRT OFI-EC	<2 a	<2 a	<2 a	<2 a	<2 a
	GR CTR	<2 a	<2 a	<2 a	<2 a	<2 a
	GR OFI-EC	<2 a	<2 a	<2 a	<2 a	<2 a
3 days	MRT CTR	<2 a	<2 a	<2 a	<2 a	<2 a
	MRT OFI-EC	<2 a	<2 a	<2 a	<2 a	<2 a
	GR CTR	<2 a	<2 a	<2 a	<2 a	<2 a
	GR OFI-EC	<2 a	<2 a	<2 a	<2 a	<2 a
5 days	MRT CTR	<2 a	<2 a	<2 a	<2 a	<2 a
	MRT OFI-EC	<2 a	<2 a	<2 a	<2 a	<2 a
	GR CTR	<2 a	<2 a	<2 a	<2 a	<2 a
	GR OFI-EC	<2 a	<2 a	<2 a	<2 a	<2 a
7 days	MRT CTR	3.06 ± 0.15 a	3.11 ± 0.20 a	3.04 ± 0.14 a	3.01 ± 0.25 a	2.99 ± 0.17 a
	MRT OFI-EC	2.14 ± 0.13 b	2.25 ± 0.19 b	2.17 ± 0.13 b	2.20 ± 0.12 b	2.15 ± 0.23 b
	GR CTR	3.24 ± 0.24 a	3.35 ± 0.21 a	3.09 ± 0.23 a	2.89 ± 0.15 a	3.11 ± 0.11 a
	GR OFI-EC	2.33 ± 0.20 b	2.15 ± 0.11 b	2.19 ± 0.10 b	2.30 ± 0.17 b	2.24 ± 0.15 b
10 days	MRT CTR	4.24 ± 0.32 a	3.88 ± 0.20 a	3.70 ± 0.25 a	4.44 ± 0.25 a	4.16 ± 0.27 a
	MRT OFI-EC	3.41 ± 0.17 b	3.01 ± 0.15 b	3.11 ± 0.19 b	3.39 ± 0.19 b	3.27 ± 0.23 b
	GR CTR	4.60 ± 0.20 a	4.16 ± 0.31 a	4.05 ± 0.21 a	4.39 ± 0.20 a	4.27 ± 0.11 a
	GR OFI-EC	3.77 ± 0.12 b	3.21 ± 0.13 b	3.40 ± 0.24 b	3.27 ± 0.14 b	3.33 ± 0.31 b
13 days	MRT CTR	5.48 ± 0.27 a	5.04 ± 0.21 a	5.12 ± 0.32 a	5.60 ± 0.19 a	5.41 ± 0.23 a
	MRT OFI-EC	4.40 ± 0.30 b	4.24 ± 0.15 b	4.05 ± 0.22 b	4.37 ± 0.20 b	4.56 ± 0.15 b
	GR CTR	5.11 ± 0.23 a	5.27 ± 0.19 a	5.01 ± 0.31 a	5.33 ± 0.25 a	5.39 ± 0.32 a
	GR OFI-EC	4.22 ± 0.19 b	4.19 ± 0.24 b	4.33 ± 0.15 b	4.49 ± 0.17 b	4.20 ± 0.19 b


Units are log CFU/g. Results indicate mean values ± S.D. of three plate counts. Data within a column followed by the same letter are not significantly different according to Tukey's test. Abbreviations: MRT CTR, uncoated fruit from the cv. *Martorana*; MRT OFI-EC, fruits from the cv. *Martorana* coated with *O. ficus-indica* mucilage; GR CTR, uncoated fruit from the cv. *Gigante Rossa*; GR OFI-EC, fruits from the cv. *Gigante Rossa* coated with *O. ficus-indica* mucilage.

According to Tukey's test, statistically significant differences were found for the levels of TMM, TPM, pseudomonads, yeasts, and molds among untreated and coated loquat samples of both cultivars. These differences were observed at seven days of refrigerate storage, when all five microbial groups appeared in all trials. In particular, these microorganisms were found at 10<sup>3</sup> CFU/g in MRT CTR and GR CTR and at 10<sup>2</sup> CFU/g in MRT OFI-EC and


GR OFI-EC. The concentrations of bacteria and fungi in all productions increased during the entire cold storage period but the MRT OFI-EC and GR OFI-EC treatments showed final counts 1 log cycle lower than those estimated for control productions.

### 3.6. Sensory Analysis and Visual Score

Uncoated MRT, GR (CTR), and coated MRT and GR (OFI-EC) loquat fruits samples were subjected to sensory evaluation on each sampling date. Minimally processed loquat fruits' sensory profiles were positively affected by mucilage coating; indeed, panelists preferred MRT OFI-EC and GR OFI-EC samples on each sampling date with mean scores 1.2 times higher than those of MRT CTR and GR CTR during the cold storage period (Figures 6 and 7). Additionally, at the end of the storage period (13 days), OFI-EC samples were preferred by judges, showing the highest scores in almost all sensorial parameters. MRT OFI-EC and GR OFI-EC samples showed mean scores 1.5 and 1.3 times higher, respectively, in terms of sensory evaluation than those of MRT CTR and GR CTR samples (Figures 6 and 7).



**Figure 6.** Sensorial analysis of minimally processed *Martorana* loquat fruit, untreated fruit (MRT CTR), and fruit treated with mucilage (MRT OFI-EC) after 3 (T3), 5 (T5), 7 (T7), 10 (T10), and 13 (T13) days of cold storage at 5 °C. Data are the mean  $\pm$  SE ( $n = 5$ ).



**Figure 7.** Sensorial analysis of minimally processed *Gigante Rossa* loquat fruit, untreated fruit (GR CTR), and fruits treated with mucilage (GR OFI-EC) after 3 (T3), 5 (T5), 7 (T7), 10 (T10), and 13 (T13) days of cold storage at 5 °C. Data are the mean  $\pm$  SE ( $n = 5$ ).


In particular, panelists perceived the largest difference in taste and texture descriptors in MRT OFI-EC samples with scores about 2 times higher than those of MRT CTR ones, and in the off-flavor descriptor, with scores 4 times lower than those of MRT CTR ones at the end of the cold storage period (Figure 6). Panelists perceived the largest difference in sweetness and aroma descriptors in GR OFI-EC samples, with scores more than 2 times higher than those of GR CTR ones, and in the off-flavor descriptor, with scores 5 times lower than those of MRT CTR ones at the end of the cold storage period (Figure 7).

Panelists perceived off flavor in MRT CTR and in GR CTR samples from days 5 to 13 at 5 °C, while the perception of this descriptor was almost absent in MRT OFI-EC and GR OFI-EC samples on each sampling date (Figures 6 and 7).


The sensory analysis showed that judges had a higher preference for coated samples at the end of the cold storage period. The mucilage coating did not negatively affect the natural taste of loquat fruits, which is an important aspect regarding the use of edible coatings when taste modification is undesirable; indeed, OFI edible coating exalted some important parameters, as well as aroma, sweetness, and taste, which are particularly appreciated by consumers.



The visual score of MRT CTR and GR CTR samples significantly decreased during storage; indeed, CTR samples of both cultivars had a severe descending trend that dropped below the limit of marketability and edibility after 7 days and 10 days of storage, respectively (Figures 8 and 9). In contrast, the MRT OFI-EC samples recorded visual scores above the limit of marketability until 10 days of cold storage and edibility until the end of the cold storage period (Figure 8). *Gigante Rossa* loquat fruits showed better performance in terms of visual scores than *Martorana* ones; indeed, GR OFI-EC recorded visual scores above the limit of marketability and edibility until the end of the cold storage period (Figure 8).



**Figure 8.** Visual score of minimally processed *Martorana* loquat fruits, untreated fruit (MRT CTR), and fruits treated with mucilage (MRT OFI-EC) during cold storage (13 days at 5 °C). Different lowercase letters indicate significant differences between the treatments on each sampling date. Data are the mean ± SE (vertical bars represent standard error;  $n = 5$ ). Data are the mean ± SE (vertical bars represent standard error;  $n = 5$ ) with 5 = very good, 4 = good, 3 = fair (limit of marketability), 2 = poor (limit of edibility) and 1 = very poor (inedible).



**Figure 9.** Visual score of minimally processed *Gigante Rossa* loquat fruits, untreated fruit (GR CTR), and fruits treated with mucilage (GR OFI-EC) during cold storage (13 days at 5 °C). Different lowercase

letters indicate significant differences between the treatments on each sampling date. Data are the mean  $\pm$  SE (vertical bars represent standard error;  $n = 5$ ). Data are the mean  $\pm$  SE (vertical bars represent standard error;  $n = 5$ ) with 5 = very good, 4 = good, 3 = fair (limit of marketability), 2 = poor (limit of edibility) and 1 = very poor (inedible).

MRT OFI-EC and GR OFI-EC samples showed visual scores 5 and 6 times higher than those of MRT CTR and GR CTR ones, respectively, at the end of the cold storage period (Figures 8 and 9), confirming that the OFI edible coating positively affects the overall loquat fruit appearance.

#### 4. Discussion

Loquat fruits are a non-climacteric fruit, with a short postharvest life due to an intrinsic predisposition to physical damage and microbial decay. The fruit decays quickly after harvest, and losses in firmness, taste, and juiciness occur rapidly during postharvest life [1].

Loquat fruit are easily bruised and scratched, and the damaged areas usually later turn brown or black in air. In addition, low temperature injury is one of the major limitations for long-term cold storage of loquat fruit. Fruit processing as well as peeling could allow producers to also sell to the market loquat fruits that present large purple spotted areas in the epicarp, which are usually considered unmarketable.

Fruit texture is an important quality attribute in minimally processed fruits, as the enzymatic reactions due to fruit processing operations (peeling, slicing, etc.) lead to rapid losses in firmness [24]. In our study, the highest fruit firmness values were measured in OFI coated samples in both cultivars during the cold storage period, showing the ability of mucilage to preserve fruit structure (Table 1). This effect on fruit firmness could be attributed to calcium content in *Opuntia ficus-indica* (OFI) mucilage, which preserves the fruit cell wall integrity by interacting with the pectic acid in the cell walls to form calcium pectate [25]. Previous studies reported that fruit calcium pre-harvest and postharvest treatments increased calcium content in the fruit, maintaining firmness in strawberry [13], fig [26], guava fruits [27], peach [28], and cactus pear fruits [8].

Our study showed the positive effect of polysaccharidic coatings, such as cactus pear mucilage, that act as a barrier reducing losses in firmness, as reported in previous studies [8]. Indeed, MRT OFI-EC and GR OFI-EC samples reported firmness values 1.1 times higher than MRT CTR and GR CTR samples at end of the cold storage period, enhancing their resistance to mechanical damage during storage and, thereby, reducing economic losses during the food chain.

TSS, TA, extractable juice, and ascorbic acid content are important indicators to measure the quality of loquat fruit; the changes in composition and content affect the fruit's taste and acceptance. Concerning chemical parameters, there was evidence of slight changes in terms of TSS and TA during storage in both MRT and GR CTR and MRT and GR OFI-EC samples (Table 1). OFI-EC edible coating inhibited the decrease of TSS, TA, extractable juice, and ascorbic acid content in both cultivars; the edible coating treatment, in terms of TSS content, was more effective in *Gigante Rossa* loquat fruit, showing higher values on all sampling dates. In contrast, the ascorbic acid content in *Martorana* loquat fruits was higher than that in *Gigante Rossa* ones. The effect of OFI-EC edible coating on limiting the reduction of TSS and juiciness loss was probably related to reducing respiration, transpiration, and metabolic activity, thereby retarding the senescence process of loquat fruit, in a similar way to that reported by Wang et al. [4] with nano-SiO<sub>2</sub> packing treatment.

One of the most beneficial effects of edible coating is the maintenance of high RH inside the fruit packaging, with OFI mucilage acting as a barrier to water transfer, reducing the weight loss [10]. The results of our study showed that the weight loss of minimally processed loquat fruit was positively affected by OFI mucilage coating; indeed, MRT and GR coated samples showed weight loss values 1.5 times higher than MRT and GR uncoated ones (Figures 1 and 2). OFI edible coating was significantly effective in preventing weight

loss in both cultivars, as reported by previous studies on strawberry [7,13], kiwifruit [29], and cactus pear [8].

Color is one of the main factors that affect consumer fruit choice and acceptability, and the pulp of loquat fruit contains carotenoids, vitamins B1 and B2 and nicotinamide [2].

Carotenoids are mainly responsible for the color of loquat fruit, the most abundant ones being  $\beta$ -carotene,  $\beta$ -cryptoxanthin, lutein, violaxanthin,  $\alpha$ -carotene, and  $\gamma$ -carotene [30]. Flesh browning is a serious problem for postharvest storage and processing of loquat fruit; the fruit turns brown rapidly when peeled or crushed. During storage, fruit browning occurs from the core area and is accompanied by lignification of the flesh tissue [2].

In our study, MRT and GR uncoated loquat fruits' brightness decreased significantly during cold storage, while MRT and GR coated samples showed slight change in brightness during cold storage with values 1.4 and 1.8 times higher than those of MRT and GR uncoated ones at the end of the cold storage period (Figure 3). Fruit color decrease and browning are mainly caused by enzymatic oxidation of endogenous polyphenols into quinones, which are then polymerized with other quinones and amines to form brown pigments [2].

OFI mucilage edible coating has the potential to act as an effective barrier against gaseous exchange between the environment and coated fruit by reducing  $O_2$  permeability and promoting  $CO_2$  accumulation in the atmosphere around the fruit [10].

In-package  $CO_2$  and  $O_2$  concentration increased and decreased (Figure 4A,B), respectively, during the cold storage period in both treatments (CTR and OFI-EC). In-package atmosphere values fluctuated between 0.04 and 12.07 kPa, and 0.04 and 8.20 kPa for  $CO_2$  in MRT and GR uncoated loquat fruits, respectively and between 20 and 6.07 kPa, and 20 and 7.11 kPa in MRT and GR uncoated loquat fruits, respectively, for  $O_2$  from the beginning to the end of the cold storage period (Figure 4A,B). In-package atmosphere values fluctuated between 0.04 and 5.13 kPa, and 0.04 and 4.00 kPa for  $CO_2$  in MRT and GR coated loquat fruits, respectively, and between 20 and 9.27 kPa, and 20 and 11.23 kPa in MRT and GR coated loquat fruits, respectively, for  $O_2$  from the beginning to the end of the cold storage period (Figure 4A,B). Differences among treatments (CTR and OFI-EC) were consistent during storage, with significant reduction of the respiration rate of coated loquat fruits in both treatments, confirming the gas barrier properties of OFI mucilage on the fruits, as reported elsewhere [8,31].

Several studies showed that phenolic compounds acted as non-enzymatic antioxidant and their accumulation might play a strong role in free radical scavenging and keeping cells from the oxidative damage caused by free radicals in fruits and vegetables [4,32]. The antioxidant capacity after the processing operations could be increased by some factors (i.e., phenols, betalains, vitamin C) and decreased by others, and its trend would reflect the contribution given by each individual factor [24].

Our study showed that OFI edible coating treatment increased total phenolic content in both cultivars which, accompanied with higher levels of DPPH and hydroxyl radical scavenging rate in MRT and GR coated loquat fruits (Table 2), reduced the damage of reactive oxygen species (ROS). Our results confirmed that total phenolic content was positively correlated with DPPH and hydroxyl radical scavenging capacities, and negatively correlated with the ROS damage in loquat fruit, as reported by several other studies [4,32]. OFI edible coating treatment did not affect the total carotenoids content in either cultivar.

Furthermore, *Martorana* loquat fruits showed higher total phenolic content than *Gigante Rossa* ones, contributing to higher DPPH and hydroxyl radical scavenging capacities, which could be related to the varietal and genetic difference. The significant difference in terms of antioxidant capacity between white-flesh and orange-red-flesh loquat cultivars was also observed in other studies [4,5,33]. Our study showed that OFI edible coating, with the increase of total phenolic content, could play a positive role in scavenging reactive oxygen species, contributing to protecting cell membrane peroxidation and damage, therewith delaying the loquat fruit senescence.

Loquat is a very perishable fruit, and its shelf life is usually influenced by postharvest decay determined by bacteria, yeasts, and molds [2,34]. For this reason, as with all minimally processed fruits and vegetables, the evaluation of microbial composition of loquat fruits deserves particular attention [35]. In this study, *O. ficus-indica* mucilage was characterized for antibacterial activity and microbial load before edible coating application on loquat fruits. The inhibitory activity was determined against spoilage and pathogenic bacteria commonly associated with foods of plant origin [36–38]. Only *E. coli* was resistant to the *O. ficus-indica* mucilage edible coating assayed. Interestingly, the coating strongly inhibited *Pseudomonas*, which are responsible for the alteration of different vegetables and fruits [39], and *L. monocytogenes*, *S. enteritidis*, and *St. aureus*, which are responsible for outbreaks associated with vegetables consumed raw [40]. Neither bacteria nor fungi developed from the serial dilutions of the *O. ficus-indica* mucilage analyzed. The absence of microorganisms at detectable levels in *O. ficus-indica* mucilage has already been reported by Liguori et al. [7,8]. None of the loquat fruit samples analyzed revealed the presence of pathogenic bacteria. The same trend was reported by Allegra et al. [29] in kiwifruit slices coated with *O. ficus-indica* mucilage as edible coating. TMM, TPM, pseudomonads, yeasts, and molds in control and treated fruits appeared at seven days of refrigerate storage and increased over time. This trend was previously reported by other authors for loquat fruits uncoated and treated with edible films extracted from cherry and apricot tree gums [41]. Although bacteria and fungi increased during storage in coated loquat fruits, cell densities observed in untreated fruits were 1 log cycle higher. This trend is mainly due to the ability of *O. ficus-indica* mucilage edible coatings to reduce the microbial growth in fresh-cut fruits [42].

The sensory analysis showed that judges preferred mucilage coated samples at each sampling date of the cold storage period, as reported by previous studies on other fruits [7,8,26,29]. MRT OFI-EC and GR OFI-EC loquat fruits samples were preferred by the panelist in all the descriptors that gave mean scores of 7.4 and 8.0 to overall acceptance, respectively, at the end of the cold storage period (5 °C), while MRT CTR and GR CTR had mean scores of 6.2 and 6.8 in overall acceptance, respectively, at the end of the cold storage period (5 °C) (Figures 6 and 7). Panelists gave scores 1.6 times higher in terms of overall acceptance to MRT OFI-EC and GR OFI-EC loquat fruits samples, respectively, than MRT CTR and GR CTR ones, at the end of the cold storage period (5 °C). The mucilage coating did not negatively affect the natural taste of loquat fruits, which is an important aspect regarding the use of edible coatings when taste modification is undesirable; indeed, OFI edible coating exalted some important parameters, as well as aroma, sweetness, and taste, that are particularly appreciated by consumers.

OFI-EC loquat fruits had the highest visual quality scores until the end of the cold storage period, while MRT OFI-EC samples were above the limit of marketability until 10 days of cold storage and edibility until the end of the cold storage period; GR OFI-EC samples were still above the marketability and edibility threshold during the storage period, while MRT CTR and GR CTR samples were marketable until 3 days of cold storage and edible until the first 6 days of storage (Figures 8 and 9). As reported by previous studies [7,8,29], mucilage coating positively affects the overall appearance of loquat during cold storage, which had a severe descending trend in uncoated loquat fruits.

## 5. Conclusions

The aim of our study was to assess the effects of *O. ficus-indica* mucilage-based coating on quality, nutraceutical value, microbiological growth, and sensorial parameters of minimally processed white-flesh *Martorana* and orange-red-flesh *Gigante Rossa* loquat fruits during cold storage.

Our data showed a significant effect of mucilage coating on preserving quality, nutraceutical value, sensorial parameters, and improving postharvest life of minimally processed loquat fruits. *O. ficus-indica* mucilage had a barrier effect on loquat minimally processed fruit during cold storage, reflected by coated samples having lower weight loss,

higher firmness, and a lower respiration rate than uncoated ones, after 13 days of cold storage at 5 °C. This factor could reduce economic losses; loquat fruit are easily bruised and scratched, and the damaged areas usually later turn brown or black in air.

OFI-EC edible coating inhibited the decrease of TSS, TA, extractable juice, and ascorbic acid content in both cultivars; the edible coating treatment, in terms of TSS content, was more effective in *Gigante Rossa* loquat fruit, showing higher values on all sampling dates. In contrast, the ascorbic acid content in *Martorana* loquat fruits was higher than that in *Gigante Rossa* ones.

Furthermore, OFI edible coating, with an increase of total phenolic content, could play a positive role in scavenging reactive oxygen species, contributing to protecting cell membrane peroxidation and damage, therewith delaying the loquat fruit senescence.

The application of edible coating of *O. ficus-indica* mucilage was not able to inhibit microbial growth below the detection limit during the entire period of analysis but reduced significantly their development in coated loquat fruits of *Martorana* and *Gigante Rossa* cultivars.

Visual quality and sensorial analysis showed that judges had a higher preference for coated loquat fruits than uncoated ones at the end of the cold storage period. Furthermore, mucilage coating did not negatively affect the natural taste of loquat fruits, which is an important aspect regarding the use of edible coatings when taste modification is undesirable. Indeed, mucilage coating exalted some important parameters, as well as firmness, brightness, aroma, sweetness, and taste, which are particularly appreciated by consumers.

Our study suggested that minimally fruit processing, as well as coating with *O. ficus-indica* mucilage, could improve peeled loquat fruits' shelf life and allow producers to also sell to the market loquat fruits that present large spotted areas in the epicarp, which are usually considered unmarketable.

**Author Contributions:** Conceptualization, G.L.; methodology, G.L., G.G. and R.G.; validation, G.L., R.G., P.I. and L.S.; formal analysis, G.L., G.G., R.G. and A.A.; investigation, G.L., G.G. and R.G.; resources, G.L. and L.S.; data curation, G.L. and R.G.; writing—original draft preparation, G.L., G.G., R.G., L.S. and P.I.; writing—review and editing, G.L., G.G., R.G., L.S., A.A. and P.I.; visualization, G.L., G.G., R.G., L.S., A.A. and P.I.; supervision, G.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Liguori, G.; Farina, V.; Sortino, G.; Mazzaglia, A.; Inglese, P. Effects of 1-methylcyclopropene on postharvest quality of white- and yellow-flesh loquat (*Eriobotrya japonica* Lindl.) fruit. *Fruits* **2014**, *69*, 363–370. [[CrossRef](#)]
2. Pareek, S.; Benkeblia, N.; Janick, J.; Cao, S.; Yahia, E.M. Postharvest physiology and technology of loquat (*Eriobotrya japonica* Lindl.) fruit. *J. Sci. Food Agric.* **2014**, *94*, 1495–1504. [[CrossRef](#)]
3. Petriccione, M.; Pasquariello, M.S.; Mastrobuni, F.; Zampella, L.; Di Patre, D.; Scortichini, M. Influence of a chitosan coating on the quality and nutraceutical traits of loquat fruit during postharvest life. *Sci. Hort.* **2015**, *197*, 287–296. [[CrossRef](#)]
4. Wang, L.; Shao, S.; Madebo, M.P.; Hou, Y.; Zheng, Y.; Jin, P. Effect of nano-SiO<sub>2</sub> packing on postharvest quality and antioxidant capacity of loquat fruit under ambient temperature storage. *Food Chem.* **2020**, *315*, 126295. [[CrossRef](#)] [[PubMed](#)]
5. Cao, S.F.; Yang, Z.F.; Cai, Y.T.; Zheng, Y.H. Fatty acid composition and antioxidant system in relation to susceptibility of loquat fruit to chilling injury. *Food Chem.* **2011**, *127*, 1777–1783. [[CrossRef](#)]
6. Raffo, A.; Paoletti, F. Fresh-Cut Vegetables Processing: Environmental Sustainability and Food Safety Issues in a Comprehensive Perspective. *Front. Sustain. Food Syst.* **2022**, *5*, 681459. [[CrossRef](#)]
7. Liguori, G.; Gaglio, R.; Settanni, L.; Inglese, P.; D'Anna, F.; Miceli, A. Effect of *Opuntia ficus-indica* Mucilage Edible Coating in Combination with Ascorbic Acid, on Strawberry Fruit Quality during Cold Storage. *J. Food Qual.* **2021**, *2021*, 9976052. [[CrossRef](#)]
8. Liguori, G.; Gaglio, R.; Greco, G.; Gentile, C.; Settanni, L.; Inglese, P. Effect of *Opuntia ficus-indica* Mucilage Edible Coating on Quality, Nutraceutical, and Sensorial Parameters of Minimally Processed Cactus Pear Fruits. *Agronomy* **2021**, *11*, 1963. [[CrossRef](#)]
9. Allegra, A.; Gallotta, A.; Carimi, F.; Mercati, F.; Inglese, P.; Martinelli, F. Metabolic profiling and post-harvest behavior of “Dottato” Fig (*Ficus carica* L.) fruit covered with an edible coating from *O. ficus-indica*. *Front. Plant Sci.* **2018**, *9*, 1321. [[CrossRef](#)] [[PubMed](#)]
10. Gheribi, R.; Khwaldia, K. Cactus Mucilage for Food Packaging Applications. *Coatings* **2019**, *9*, 655. [[CrossRef](#)]
11. GirmaAbera, N.; Kebede, W.; Wassu, M. Effect of Aloe gel and cactus mucilage coating on chemical quality and sensory attributes of mango (*Mangifera indica* L.). *J. Postharvest Technol.* **2019**, *7*, 31–43.

12. Sogvar, O.B.; Saba, M.K.; Emamifar, A. Aloe vera and ascorbic acid coatings maintain postharvest quality and reduce microbial load of strawberry fruit. *Postharvest Biol. Technol.* **2016**, *114*, 29–35. [[CrossRef](#)]
13. Del Valle, V.; Hernández-Muñoz, P.; Guarda, A.; Galotto, M.J. Development of a cactus-mucilage edible coating (*Opuntia ficus indica*) and its application to extend strawberry (*Fragaria ananassa*) shelf-life. *Food Chem.* **2005**, *91*, 751–756. [[CrossRef](#)]
14. Riaz, S.; Sultan, M.T.; Sibte-e-Abass, M.; Shabir Ahmad, M.I.R.; Bilal Hussain, M.; Shariati, M.A.; Kosenko, I.S.; Kleymenova, N.L.; Egorova, G.N. Extraction of polysaccharides from *Opuntia* cactus for its potential application in edible coating to improve the shelf life of citrus (Kinnow Mandarin) fruit. *J. Microbiol. Biotechnol. Food Sci.* **2018**, *8*, 745–750. [[CrossRef](#)]
15. Sortino, G.; Allegra, A.; Farina, V.; De Chiara, M.L.V.; Inglese, P. Genotype influence on shelf life behaviour of minimal processed loquat (*Eriobotrya japonica* (Thunb.) Lindl.) fruit: The role of sugar, acid organics and phenolic compounds. *Chem. Biol. Technol. Agric.* **2022**, *9*, 8. [[CrossRef](#)]
16. Du Toit, A.; De Wit, M. A Process for Extracting Mucilage from *Opuntia ficus-indica* and *Aloe barbadensis*. South. Africa Patent No. PA153178/P, 12 May 2011.
17. Passafiume, R.; Gugliuzza, G.; Gaglio, R.; Busetta, G.; Tinebra, I.; Sortino, G.; Farina, V. Aloe-Based Edible Coating to Maintain Quality of Fresh-Cut Italian Pears (*Pyrus communis* L.) during Cold Storage. *Horticulturae* **2021**, *7*, 581. [[CrossRef](#)]
18. Kanner, J.; Harel, S.; Granit, R. Betalains-A new class of dietary cationized antioxidants. *J. Agric. Food Chem.* **2001**, *49*, 5178–5185. [[CrossRef](#)]
19. Singleton, V.; Rossi, J. Colorimetry of total polyphenols with phospho-molybdic phosphor tungstic acid reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144–158.
20. Kichtenthaler, H.K.; Wellburn, A.R. Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvent. *Biochem. Soc. Trans.* **1983**, *603*, 591–593. [[CrossRef](#)]
21. Ranganna, S. *Manual of Analysis of Fruit and Vegetable Products*; McGraw-Hill Publishing Co. Ltd.: New Delhi, India, 1977.
22. Gaglio, R.; Barbera, M.; Aleo, A.; Lommatzsch, I.; La Mantia, T.; Settanni, L. Inhibitory activity and chemical characterization of *Daucus carota* subsp. *maximus* essential oils. *Chem. Biod.* **2017**, *14*, e1600477.
23. Miceli, A.; Aleo, A.; Corona, O.; Sardina, M.T.; Mammìna, C.; Settanni, L. Antibacterial activity of *Borago officinalis* and *Brassica juncea* aqueous extracts evaluated in vitro and in situ using different food model systems. *Food Control* **2014**, *40*, 157–164. [[CrossRef](#)]
24. Palma, A.; Continella, A.; La Malfa, S.; D’Aquino, S. Changes in physiological and some nutritional, nutraceuticals, chemical-physical, microbiological and sensory quality of minimally processed cactus pears cvs ‘Bianca’, ‘Gialla’ and ‘Rossa’ stored under passive modified atmosphere. *J. Sci. Food Agric.* **2018**, *98*, 1839–1849. [[CrossRef](#)] [[PubMed](#)]
25. Irfan, P.K.; Vanjakshi, V.; Keshava Prakash, M.N.; Ravi, R.; Kudachikar, V.B. Calcium chloride extends the keeping quality of fig fruit (*Ficus carica* L.) during storage and shelf-life. *Postharvest Biol. Technol.* **2013**, *82*, 70–75. [[CrossRef](#)]
26. Allegra, A.; Sortino, G.; Inglese, P.; Settanni, L.; Todaro, A.; Gallotta, A. The effectiveness of *Opuntia ficus-indica* mucilage edible coating on post-harvest maintenance of ‘Dottato’ fig (*Ficus carica* L.) fruit. *Food Packag. Shelf Life* **2017**, *12*, 135–141. [[CrossRef](#)]
27. Singh Bhooriya, M.; Bisen, B.P.; Pandey, S.K. Effect of post-harvest treatments on shelf life and quality of Guava (*Psidium guajava*) fruits. *Int. J. Chem. Stud.* **2018**, *6*, 2559–2564.
28. Manganaris, G.A.; Vasilakakis, M.; Diamantidis, G.; Mignani, I. The effect of postharvest calcium application on tissue calcium concentration, quality attributes, incidence of flesh browning and cell wall physicochemical aspects of peach fruits. *Food Chem.* **2007**, *100*, 1385–1392. [[CrossRef](#)]
29. Allegra, A.; Inglese, P.; Sortino, G.; Settanni, L.; Todaro, A.; Liguori, G. The influence of *Opuntia ficus-indica* mucilage edible coating on the quality of ‘Hayward’ kiwifruit slices. *Postharvest Biol. Technol.* **2016**, *120*, 45–51. [[CrossRef](#)]
30. Ding, C.K.; Chachin, K.; Hamazu, Y.; Ueda, Y.; Imahori, Y. Effects of storage temperatures on physiology and quality of loquat fruit. *Postharvest Biol. Technol.* **1998**, *14*, 309–315. [[CrossRef](#)]
31. Christopoulos, M.V.; Gkatzos, D.; Kafkaletou, M.; Bai, J.; Fanourakis, D.; Tsaniklidis, G.; Tsantili, E. Edible Coatings from *Opuntia ficus-indica* Cladodes Alongside Chitosan on Quality and Antioxidants in Cherries during Storage. *Foods* **2022**, *11*, 699. [[CrossRef](#)]
32. Xu, H.X.; Chen, J.W.; Xie, M. Effect of different light transmittance paper bags on fruit quality and antioxidant capacity in loquat. *J. Sci. Food Agric.* **2010**, *90*, 1783–1788. [[CrossRef](#)]
33. Yang, S.L.; Sun, C.D.; Wang, P.; Shan, L.L.; Cai, C.; Zhang, B.; Zhang, W.; Ferguson, I.; Chen, K.S. Expression of expansin genes during postharvest lignification and softening of ‘Luoyangqing’ and ‘Baisha’ loquat fruit under different storage conditions. *Postharvest Biol. Technol.* **2008**, *49*, 46–53. [[CrossRef](#)]
34. Yang, H.; Wang, L.; Li, S.; Gao, X.; Wu, N.; Zhao, Y.; Sun, W. Control of postharvest grey spot rot of loquat fruit with *Metschnikowia pulcherrima* E1 and potential mechanisms of action. *Biol. Control* **2021**, *152*, 104406. [[CrossRef](#)]
35. Brackett, R. Microbiological consequences of minimally processed fruits and vegetables. *J. Food Qual.* **1987**, *10*, 195–206. [[CrossRef](#)]
36. Miceli, A.; Gaglio, R.; Francesca, N.; Ciminata, A.; Moschetti, G.; Settanni, L. Evolution of shelf-life parameters of ready-to-eat escarole (*Cichorium endivia* var. *latifolium*) subjected to different cutting operations. *Sci. Hort.* **2019**, *247*, 175–183.
37. European Food Safety Authority. Scientific opinion on the risk posed by pathogens in food of non-animal origin. Part 1. *EFSA J.* **2013**, *11*, 3025. [[CrossRef](#)]
38. Miceli, A.; Settanni, L. Influence of agronomic practices and pre-harvest conditions on the attachment and development of *Listeria monocytogenes* in vegetables. *Ann. Microbiol.* **2019**, *69*, 185–199. [[CrossRef](#)]

39. Barth, M.; Hankinson, T.R.; Zhuang, H.; Breidt, F. Microbiological Spoilage of Fruits and Vegetables. In *Compendium of the Microbiological Spoilage of Foods and Beverages*; Food Microbiology and Food Safety; Sperber, W.H., Doyle, M.P., Eds.; Springer: New York, NY, USA, 2009; pp. 135–183.
40. Leff, J.W.; Fierer, N. Bacterial communities associated with the surfaces of fresh fruits and vegetables. *PLoS ONE* **2013**, *8*, e59310. [[CrossRef](#)]
41. Ergin, S.Ö.; Yaman, H.; Dilek, M. The usage of edible films extracted from cherry and apricot tree gums for coating of strawberry (*Fragaria ananassa*) and loquat (*Eriobotrya japonica*) fruits. *Turk. J. Agric. Food Sci. Technol.* **2018**, *6*, 561–569.
42. Barba, F.J.; Garcia, C.; Fessard, A.; Munekata, P.E.; Lorenzo, J.M.; Aboudia, A.; Ouadia, A.; Remize, F. *Opuntia ficus indica* edible parts: A food and nutritional security perspective. *Food Rev. Int.* **2022**, *38*, 930–952. [[CrossRef](#)]